## **References:** applications to wireless communications

- Gray R. M. On the asymptotic eigenvalue distribution of Toeplitz matrices. IEEE Trans. Inform. Theory 18 (1972), 725–730.
- [2] Müller, R. A random matrix model of communication via antenna arrays. IEEE Trans. Inform. Theory 48 (2002), no. 9, 2495–2506.
- [3] Özgür, A., Lévêque, Tse D. Hierarchical cooperation achieves optimal capacity scaling in ad hoc networks. Preprint.
- [4] Telatar, I. E. Capacity of multi-antenna Gaussian channels. European Trans. on Telecom. 10 (1999), no. 6, 585–595.
- [5] Telatar, I. E., Tse, D. N. C. Capacity and mutual information of wideband multipath fading channels. IEEE Trans. Inform. Theory 46 (2000), no. 4, 1384–1400.
- [6] Tse, D. N. C., Hanly, S. V. Linear multiuser receivers: effective interference, effective bandwidth and user capacity. IEEE Trans. Inform. Theory 45 (1999), no. 2, 641–657.
- [7] Tse D. N. C., Zeitouni O. Linear multiuser receivers in random environments. IEEE Trans. Inform. Theory 46 (2000), no. 1, 171–188.
- [8] Tulino, A. M., Verdú, S. Random matrix theory and wireless communications. Foundations and Trends in Communications and Information Theory 1 (1), Now Publishers Inc., 2004.
- [9] Verdú, S. Spectral efficiency in the wideband regime. Special issue on Shannon theory: perspective, trends, and applications. IEEE Trans. Inform. Theory 48 (2002), no. 6, 1319–1343.
- [10] Zheng, L., Tse, D. N. C. Communication on the Grassmann manifold: a geometric approach to the noncoherent multiple-antenna channel. IEEE Trans. Inform. Theory 48 (2002), no. 2, 359–383.
- [11] Zheng, L., Tse, D. N. C. Diversity and multiplexing: a fundamental tradeoff in multiple antenna channels. IEEE Transactions on Information Theory, vol. 49 (2003), 1073–1096.