

ÉCOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE

School of Computer and Communication Sciences

Handout 2
Homework 1

Information Theory and Coding
Sep. 18, 2012

PROBLEM 1. Three events E_1 , E_2 and E_3 , defined on the same space, have probabilities $P(E_1) = P(E_2) = P(E_3) = 1/4$. Let E_0 be the event that one or more of the events E_1 , E_2 , E_3 occurs.

(a) Find $P(E_0)$ when:

- (1) The events E_1 , E_2 and E_3 are disjoint.
- (2) The events E_1 , E_2 and E_3 are statistically independent.
- (3) The events E_1 , E_2 and E_3 are in fact three names for the same event.

(b) Find the maximum value $P(E_0)$ can assume when:

- (1) Nothing is known about the independence or disjointness of E_1 , E_2 , E_3 .
- (2) It is known that E_1 , E_2 and E_3 are pairwise independent, i.e., that the probability of realizing both E_i and E_j is $P(E_i)P(E_j)$, $1 \leq i \neq j \leq 3$, but nothing is known about the probability of realizing all three events together.

PROBLEM 2. A dishonest gambler has a loaded die which turns up the number 1 with probability $2/3$ and the numbers 2 to 6 with probability $1/15$ each. Unfortunately, he has left his loaded die in a box with two honest dice and can not tell them apart. He picks one die (at random) from the box, rolls it once, and the number 1 appears. Conditional on this result, what is the probability that he picked up the loaded die? He now rolls the dice once more and it comes up 1 again. What is the possibility after this second rolling that he has picked up the loaded die?

PROBLEM 3. Suppose the random variables A , B , C , D form a Markov chain: $A \ominus B \ominus C \ominus D$.

- (a) Is $A \ominus B \ominus C$?
- (b) Is $B \ominus C \ominus D$?
- (c) Is $A \ominus (B, C) \ominus D$?
- (d) Is $A \ominus B \ominus (C, D)$?

PROBLEM 4. Suppose the random variables A , B , C , D satisfy $A \ominus B \ominus C$, and $B \ominus C \ominus D$. Does it follow from these that $A \ominus B \ominus C \ominus D$?

PROBLEM 5. Let X and Y be two random variables.

- (a) Prove that the expectation of the sum of X and Y , $E[X + Y]$, is equal to the sum of the expectations, $E[X] + E[Y]$.
- (b) Prove that if X and Y are statistically independent, then X and Y are also uncorrelated (by definition X and Y are uncorrelated if $E[XY] = E[X]E[Y]$). Find an example in which X and Y are statistically dependent yet uncorrelated.

- (c) Prove that if X and Y are statistically independent, then the variance of the sum $X + Y$ is equal to the sum of variances. Is this relationship valid if X and Y are uncorrelated but not statistically independent?

PROBLEM 6. After summer, the winter tyres of a car (with four wheels) are to be put back. However, the owner has forgotten which tyre goes to which wheel, and the tyres are installed ‘randomly’, each of the $4! = 24$ permutations being equally likely.

- (a) What is the probability that tyre 1 is installed in its original position?
- (b) What is the probability that all the tyres are installed in their original positions?
- (c) What is the expected number of tyres that are installed in their original positions?
- (d) Redo the above for a vehicle with n wheels.
- (e) (Harder.) What is the probability that none of the wheels are installed in their original positions.

PROBLEM 7. We construct an ‘inventory’ by drawing n independent samples from a distribution p . Let X_1, \dots, X_n be the random variables that represent the drawings.

Suppose X is drawn from distribution p , independent of X_1, \dots, X_n .

- (a) What is the probability that X does not appear in the inventory?
- (b) Redo (a) for the special case when p is the uniform distribution over n items.
- (c) What happens to the probability in (b) when n gets large?